



RESEARCH ARTICLE

Dietary patterns and hypertension among daily laborers at Belawan Community Health Center

Eliza Juliana Hutabarat¹, Johannes Bastira Ginting^{2*}, Pahala Maringan Jubel Simangunsong³

ABSTRACT

Background: Occupational health problems in the informal sector, particularly among daily laborers in coastal areas such as the working area of Belawan Community Health Center, require special attention. Heavy physical activity, environmental exposure, and individual characteristics are strongly suspected to be associated with fluctuations in blood pressure (systolic and diastolic) that may increase hypertension risk. This study aimed to determine the relationship between dietary patterns and hypertension among daily laborers in the working area of Belawan Community Health Center.

Method: This study uses an analytical method with a quantitative approach and a cross-sectional design to assess the relationship between dietary variables and the incidence of hypertension through a single-point data collection. This study will be conducted at the Belawan Community Health Center (Puskesmas Belawan) located at Jl. Stasiun No. 1, P. JKA Complex, with implementation starting in April and continuing through completion. The target population for this study is all daily laborers within the Belawan Community Health Center (Puskesmas Belawan) working area. The accessible population is daily laborers aged 18-49 who visited the Belawan Community Health Center during the study period and met the inclusion criteria.

The sampling technique used is purposive sampling, which determines the sample based on specific considerations and criteria tailored to the research objectives. Purposive sampling was chosen because the researcher only selected respondents who met the inclusion criteria consistent with the research objectives. Therefore, respondents were selected based on predetermined inclusion criteria and recruited during the data collection period.

Results: Univariate analysis showed that most respondents had poor dietary patterns (64.1%) and were classified as having hypertension (44.6%) or prehypertension (28.3%). The Chi-square test yielded a p-value of less than 0.001, indicating a significant relationship between dietary pattern and hypertension among daily laborers in this population. Respondents with poor dietary patterns predominantly had elevated blood pressure, while those with good dietary patterns were more likely to have normal blood pressure.

Conclusion: There is a significant relationship between dietary pattern and hypertension among daily laborers at Belawan Community Health Center. Regular health screening programs for informal sector workers in this area are recommended.

Keywords: daily laborers, dietary pattern, blood pressure

Introduction

Hypertension, or high blood pressure, is a non-communicable disease that poses a global health threat and is known as a silent killer.¹ According to WHO data, this disease, which is a primary trigger for cardiovascular complications such as heart disease and stroke, has affected more than 1.28 billion people worldwide, with the majority of cases in low- and middle-income countries. One of the major modifiable risk factors is dietary pattern.²⁻⁴ Various studies have shown that high consumption of sodium, saturated fats,

Affiliation

¹Undergraduate Program in Public Health, Universitas Prima Indonesia

²Department of Public Health, Universitas Prima Indonesia

Correspondence:

johannesbastiraginting@unprimdn.ac.id

salt, fast food, and ultra-processed foods consistently increases hypertension risk.^{5,6} Conversely, a healthy diet rich in fruits and vegetables has been proven to reduce this risk.^{7,8}

At the national level, Indonesia faces a major challenge with a hypertension prevalence of 34.1% among adults aged 18 years and above based on the 2018 Basic Health Research (Riskesdas).⁹ Vulnerability is higher among informal sector workers, with a prevalence reaching 37.5% due to stress factors and unhealthy lifestyles.¹⁰ Daily laborers, as part of the informal sector, are the most at-risk group because of economic limitations and irregular working hours.¹¹ These conditions force them to choose ready-to-eat foods or street snacks high in sodium and fat, combined with low awareness of balanced nutrition.¹² The strong relationship between poor dietary patterns and hypertension has been reinforced by research in various regions, such as at Prambanan Community Health Center, Klaten¹³ and Semanding Community Health Center, Tuban¹⁴.

A similar phenomenon occurs at the regional level in North Sumatra, particularly in Medan City. Based on the 2022 Medan City Health Profile, hypertension is among the top ten diseases, with a total of 65,904 cases. More specifically, the working area of Belawan Community Health Center recorded 1,225 active hypertension patients in 2025. Local research by Siregar et al.¹⁵ in this area confirmed that the high number of hypertension cases in Belawan is closely related to the community's habit of excessive salt consumption and low intake of fruits and vegetables. This study aimed to analyze the relationship between dietary patterns and the incidence of hypertension among daily laborers at Belawan Community Health Center.

Method

This study uses an analytical method with a quantitative approach and a cross-sectional design to assess the relationship between dietary variables and the incidence of hypertension through a single-point data collection. This study will be conducted at the Belawan Community Health Center (Puskesmas Belawan) located at Jl. Stasiun No. 1, P. JKA Complex, with implementation starting in April and continuing through completion. The target population for this study is all daily laborers within the Belawan Community Health Center (Puskesmas Belawan) working area. The accessible population is daily laborers aged 18–49 who visited the Belawan Community Health Center during the study period and met the inclusion criteria.

The sampling technique used is purposive sampling, which determines the sample based on specific considerations and criteria tailored to the research objectives. Purposive sampling was chosen because the researcher only selected respondents who met the inclusion criteria consistent with the research objectives. Therefore, respondents were selected based on predetermined inclusion criteria and recruited during the data collection period.

Results

Table 1 presents the sociodemographic characteristics of the 92 respondents. The majority were male (59 respondents, 64.1%), while female respondents numbered 33 (35.9%). Regarding age, the largest group was aged 30–35 years (40 respondents, 43.5%), followed by those aged 35–49 years (25, 27.2%), 25–30 years (17, 18.5%), and 18–25 years (10, 10.9%). In terms of education, most respondents had completed senior high school (49, 53.3%), followed by junior high school (22, 23.9%) and elementary school (21, 22.8%). All respondents (100.0%) worked as daily laborers. Regarding residential distribution, the largest group lived in Belawan I Village (38, 41.3%), with equal numbers from Belawan II Village (27, 29.3%) and Bagan Deli Village (27, 29.3%).

Table 1. Distribution of respondent characteristics (N=92)

Characteristic	Category	Frequency (n)	Percentage (%)
Sex	Male	59	64.1
	Female	33	35.9
Age	18-25 years	10	10.9
	25-30 years	17	18.5
	30-35 years	40	43.5
	35-49 years	25	27.2
Education	Elementary school	21	22.8
	Junior high school	22	23.9
	Senior high school	49	53.3
Occupation	Daily laborer	92	100.0

Village	Belawan I	38	41.3
	Belawan II	27	29.3
	Bagan Deli	27	29.3

Table 2 combines the frequency distributions for dietary pattern and blood pressure status. Regarding dietary pattern, the majority of respondents (59, 64.1%) had poor dietary patterns, characterized by frequent consumption of high-sodium foods, instant or processed foods, and fried foods. Only 33 respondents (35.9%) had good dietary patterns. Regarding blood pressure, the largest group was classified as hypertensive (41, 44.6%), followed by prehypertensive (26, 28.3%), and normal (25, 27.2%). More than two-thirds of respondents had elevated blood pressure (prehypertension or hypertension combined), indicating a substantial burden of blood pressure abnormalities in this population.

Table 2. Distribution of dietary pattern and blood pressure status (N=92)

Variable	Category	Frequency (n)	Percentage (%)
Dietary pattern	Good	33	35.9
	Poor	59	64.1
Blood pressure	Normal	25	27.2
	Prehypertension	26	28.3
	Hypertension	41	44.6

Table 3 presents the cross-tabulation between dietary pattern and blood pressure category. Among the 33 respondents with good dietary patterns, the majority had normal blood pressure (25, 27.2% of total respondents), followed by prehypertension (6, 6.5%), and only 2 (2.2%) had hypertension. In contrast, among the 59 respondents with poor dietary patterns, none had normal blood pressure. The largest group among those with poor dietary patterns had hypertension (39, 42.4%), followed by prehypertension (26, 28.3%). The Chi-square test yielded a p-value of less than 0.001, which is below the significance threshold of 0.05. Therefore, the null hypothesis is rejected, and it is concluded that there is a statistically significant relationship between dietary pattern and hypertension among daily laborers in the working area of Belawan Community Health Center. Notably, every respondent with a poor dietary pattern had elevated blood pressure (prehypertension or hypertension), while the majority of those with good dietary patterns maintained normal blood pressure.

Table 3. Relationship between dietary pattern and blood pressure (N=92)

Dietary Pattern	Normal n (%)	Prehypertension n (%)	Hypertension n (%)	Total	p-value
Good	25 (27.2)	6 (6.5)	2 (2.2)	33	<0.001
Poor	0 (0.0)	26 (28.3)	39 (42.4)	59	
Total	25 (27.2)	26 (28.3)	41 (44.6)	92	

Discussion

This study found a significant relationship between dietary pattern and hypertension among daily laborers at Belawan Community Health Center. The majority of respondents had poor dietary patterns and were in the hypertensive or prehypertensive categories. In contrast, respondents with good dietary patterns predominantly had normal blood pressure.

The high proportion of poor dietary patterns among daily laborers can be explained by their working conditions. Daily laborers typically have irregular working hours, high physical demands, and limited economic resources.¹⁶ These factors push them toward convenient, inexpensive, and calorie-dense food options such as instant noodles, fried foods, salty snacks, and processed meats available near the port area.¹⁷ Field observations and questionnaire responses indicated that respondents frequently consumed salted fish, salted eggs, dishes with excessive salt and seasoning, and instant foods.¹⁸ The coastal geography of Belawan facilitates easy access to salted and preserved fish products, and the need for savory flavors to stimulate appetite after heavy physical work further reinforces these habits.¹⁹

Physiologically, the accumulation of sodium from these foods leads to fluid retention, increased blood volume, and elevated pressure on vessel walls. This condition is worsened by high saturated fat intake from fried foods and coconut milk, which promotes plaque formation in arteries (atherosclerosis), resulting in vasoconstriction and increased peripheral resistance.²⁰ Compounding this problem is the very low awareness of nutritional balance among workers. Most respondents reported never or only occasionally consuming fruits and vegetables daily, citing inconvenience and the perception that such foods do not provide instant energy. Consequently, their bodies experience a significant deficit of potassium, magnesium, and fiber—

nutrients that help counteract sodium effects and promote blood vessel relaxation.²¹ Over time, this poor dietary pattern triggers chronic metabolic disorders such as obesity and dyslipidemia, which directly contribute to blood pressure elevation.^{21,22}

These findings are consistent with local studies by Mardianto et al.²³ and Hasanuddin et al.²⁴, which also reported significant associations between poor dietary patterns (high salt/fat, low fiber) and hypertension. The results are further supported by international evidence from Jaques et al.²¹ and a systematic review by Wang et al.²², which identified excessive sodium intake as a major cardiovascular risk factor due to increased plasma volume and cardiac workload. Moreover, the Dietary Approaches to Stop Hypertension (DASH) trial by Sacks et al.²⁵ demonstrated clinically that reducing sodium intake while increasing consumption of fruits, vegetables, and fiber-rich foods rich in potassium and magnesium is highly effective in controlling blood pressure and reducing complication risk.

This study provides specific evidence on dietary patterns and hypertension among informal sector daily laborers, a group often underrepresented in research. Direct blood pressure measurement enhanced data accuracy. However, limitations include the cross-sectional design, which cannot establish causality; the relatively small sample size (n=92) from a single health center, limiting generalizability; and reliance on self-reported dietary patterns, which may be subject to recall bias. Future studies should use larger, multi-site samples and more objective dietary assessment methods (e.g., 24-hour dietary recalls).

Belawan Community Health Center should strengthen regular health screening programs for informal sector workers, particularly daily laborers. Nutrition education interventions should focus on practical, affordable strategies to reduce sodium intake and increase fruit and vegetable consumption within the constraints of workers' schedules and budgets. Workplace-based health promotion targeting port and coastal industries could also be explored. Policy makers should consider subsidizing healthy food options for low-income workers and improving access to nutrition counseling at primary health centers.

Conclusion

This study found a significant relationship between dietary pattern and hypertension among daily laborers in the working area of Belawan Community Health Center. The majority of respondents had poor dietary patterns and were classified as having hypertension or prehypertension. Respondents with poor dietary patterns consistently had elevated blood pressure, while those with good dietary patterns were more likely to have normal blood pressure. Regular health screenings and targeted nutrition education are recommended for this high-risk population.

References

1. Mills KT, Stefanescu A, He J. The global epidemiology of hypertension. *Nat Rev Nephrol*. 2020 Apr 5;16(4):223–37.
2. Wandile PM. Hypertension and comorbidities: A silent threat to global health. *Hypertens Comorbidities*. 2024 Feb 6;1(1):1–7.
3. Sarki AM, Nduka CU, Stranges S, Kandala NB, Uthman OA. Prevalence of Hypertension in Low- and Middle-Income Countries. *Medicine (Baltimore)*. 2015 Dec;94(50):e1959.
4. Hutchinson B, Walter A, Campbell N, Whelton PK, Varghese C, Husain MJ, et al. Scaling hypertension treatment in 24 low-income and middle-income countries: economic evaluation of treatment decisions at three blood pressure cut-points. *BMJ Open*. 2024 Apr 15;14(4):e071036.
5. Wu Q, Burley G, Li L, Lin S, Shi Y. The role of dietary salt in metabolism and energy balance: Insights beyond cardiovascular disease. *Diabetes, Obes Metab*. 2023 May 28;25(5):1147–61.
6. Keller MA, Nakamura M. Acetyltransferase in cardiovascular disease and aging. *J Cardiovasc Aging*. 2024 Dec 31;4(4).
7. Belanger MJ, Kovell LC, Turkson-Ocran R, Mukamal KJ, Liu X, Appel LJ, et al. Effects of the Dietary Approaches to Stop Hypertension Diet on Change in Cardiac Biomarkers Over Time: Results From the DASH-Sodium Trial. *J Am Heart Assoc*. 2023 Jan 17;12(2).
8. Pietri P. Dietary Guidance for Cardiovascular Health: Consensus and Controversies. *Nutrients*. 2023 Oct 9;15(19):4295.
9. Kementerian Kesehatan Republik Indonesia. *Riset Kesehatan Dasar 2018 (2018 Basic Health Research)*. Jakarta; 2018.
10. Fahrurroddi DS, Tsurayya A, Santi AR, Satrya BA. Studi prevalensi kejadian hipertensi pada pekerja informal di Indonesia (Analisis data Riskesdas 2018). *Prepotif*. 2025 Dec 18;9(3):8091–7.
11. Ameya G, Ayal DY. Navigating the aftermath of the COVID-19 pandemic: unpacking food insecurity, coping strategies, and associated factors among informal workers in Addis Ababa, Ethiopia. *Cogent Food Agric*. 2024 Dec 31;10(1).
12. Prasetyono JD, Permatasari H, Setiawan A, Mulyono S, Susanto T, Mansyur M. Smoking, Nutritional Status, and Their Associations with Hypertension and Hematological Disorders Among Hotel Workers: Implications for Indonesian Occupational Health Nursing. *Int J Environ Res Public Health*. 2026 Apr 15;23(4):505.
13. Nasyaroeka AEBD, Solikhah S, Lianawati L. Food consumption pattern for hypertension patient in Klaten Regency, Central Java Province, Indonesia. *J Cakrawala Promkes*. 2024 Mar 24;6(1):1–7.
14. Mahmudah TA, Ningsih WT, Nugraheni WT, Su'udi. Pola Makan Tinggi Garam Dengan Kejadian Hipertensi Di Wilayah Kerja Puskesmas Semanding. *J Ilm Kesehat Mandira Cendikia [Internet]*. 2025 Oct 6;4(10):283–91. Available from:

- <https://journal.mandiracendikia.com/index.php/JIK-MC/article/view/1962>
15. Siregar PA, Simanjuntak SFS, Ginting FH, Tarigan S, Hanum S, Utami FS. Physical Activity, Consumption of Salty Foods and the Occurrence of Hypertension in Coastal Communities in Medan City. *J Ilm Kesehat*. 2020 Apr 30;2(1):1–8.
 16. Łokieć K, Górski-Ciebiada M. Nutritional behaviours of shift workers. *Med Ogólna i Nauk o Zdrowiu*. 2020 Sep 16;26(3):213–20.
 17. Tamba DM, Atika T, Ritonga FU. Strategi Bertahan Hidup Buruh Pengupas Kulit Udang dalam Pemenuhan Kebutuhan Dasar di Masa Covid-19. *SOSMANIORA J Ilmu Sos dan Hum*. 2023 Jun 28;2(2):278–84.
 18. Zhang X, Chen B, Jia P, Han J. Locked on salt? Excessive consumption of high-sodium foods during COVID-19 presents an underappreciated public health risk: a review. *Environ Chem Lett*. 2021 Oct 1;19(5):3583–95.
 19. Rezha DK, Hasibuan R, Maipiana DR, Lubis CM, Difhanny CN, Marpaung SY. Faktor-Faktor Risiko yang Berhubungan dengan Hipertensi di Kelurahan Belawan 1. *J Ilm Univ Batanghari Jambi*. 2023 Jul 26;23(2):1669.
 20. Kurniyanti N, Aryanti A, Makmun A, Nesyana N, Daeng Kanang IL. Hubungan Asupan Makronutrien dan Mikronutrien terhadap Tekanan Darah pada Pasien Hipertensi. *Indones J Heal*. 2022 Jun 6;2(3):147–54.
 21. Jaques DA, Wuerzner G, Ponte B. Sodium Intake as a Cardiovascular Risk Factor: A Narrative Review. *Nutrients*. 2021 Sep 12;13(9):3177.
 22. Wang YJ, Yeh TL, Shih MC, Tu YK, Chien KL. Dietary Sodium Intake and Risk of Cardiovascular Disease: A Systematic Review and Dose-Response Meta-Analysis. *Nutrients*. 2020 Sep 25;12(10):2934.
 23. Mardianto M, Darwis D, Suhartatik S. Hubungan Pola Makan Dengan Kejadian Hipertensi. *J Ilm Mhs Penelit Keperawatan [Internet]*. 2023 Nov 1;1(4). Available from: <https://ojs.institutnh.ac.id/index.php/jimpk/article/view/663>
 24. Hasanuddin A, Alwi MK, Nindrea RD, Astuti A, Dai NF, Maryam A, et al. Effect of Health Education on the Behavior of Elderly with Hypertension. *J Penelit Pendidik IPA*. 2023 Sep 25;9(9):7023–7.
 25. Sacks FM, Svetkey LP, Vollmer WM, Appel LJ, Bray GA, Harsha D, et al. Effects on Blood Pressure of Reduced Dietary Sodium and the Dietary Approaches to Stop Hypertension (DASH) Diet. *N Engl J Med*. 2001 Jan 4;344(1):3–10.
 26. Luo X, Zhang X, Ran L, Kang Z, Fu X, Mu X. Dietary sodium, potassium, and cardiometabolic risk: A cross-sectional analysis of hypertension in U.S. adults from NHANES 2017–2018. *Dang S, editor. PLoS One*. 2026 Feb 20;21(2):e0343311.